

MASSIVE STARS AND THEIR SURROUNDING NEBULAE: A COMBINED APPROACH

S. Simón-Díaz,¹ G. Stasińska,¹ J. García-Rojas,² C. Morisset,³ A. R. López-Sánchez,² and C. Esteban²

RESUMEN

Presentamos los primeros resultados de un proyecto dedicado al estudio combinado de estrellas masivas y sus regiones H II asociadas en la Galaxia. Mediante el estudio de regiones ionizadas por una única estrella pretendemos comprobar la validez de las predicciones de la nueva generación de modelos de atmósfera de estrellas masivas en términos de distribución de flujo ionizante. En nuestros análisis consideramos el efecto de las posibles distribuciones de densidad del material ionizado. Para ello hacemos uso de varios tipos de observaciones, tanto estelares como de la región H II.

ABSTRACT

We present the first results of a project aimed at the combined study of massive stars and their surrounding nebulae by means of a detailed study of Galactic H II regions ionized by only one massive star. With this, we intend to check the validity of the new generation of massive star model atmosphere codes in terms of ionizing flux distribution. We take into account the effect of the nebular density distribution in our analyses. Various types of stellar and nebular observations have been collected for this purpose.

Key Words: STARS: EARLY-TYPE — STARS: INDIVIDUAL: HD 37061 — H II REGIONS — ISM: IONIZATION MODELS — ISM: INDIVIDUAL OBJECTS: M 43

1. INTRODUCTION

The intense far ultraviolet radiation emitted by early OB-type stars ionizes the interstellar medium, generating the so-called H II regions. These ionized regions can be used to derive properties of the associated stellar population (e.g. star forming rates, age). However, since the properties of H II regions crucially depend on the spectral energy distribution (SED) of the massive star population, and this part of the stellar flux is generally unaccessible to direct observations, the predictions resulting from massive star atmosphere codes are a crucial ingredient.

The new generation of NLTE, line blanketed model atmosphere codes (Hubeny & Lanz, 1995, Hillier & Miller 1998, Pauldrach et al. 2001, Puls et al. 2005), which include a more realistic description of the physical processes characterizing the stellar atmosphere, produce quite different ionizing SEDs than the previous plane-parallel, NLTE/LTE, hydrostatic models (Mihalas & Auer 1970, Kurucz 1992, Kunze 1994). Some notes on this, and on the consequences on the ionization structure of H II regions, can be found in Gabler et al. (1989), Rubin et al. (1995), Najarro et al. (1996), Sellmaier et al.

(1996) and Stasińska & Schaerer (1997). Although new predictions seem to go in the right direction (viz. Giveon et al. 2002, Morisset et al. 2004), non-negligible differences can be found between the different stellar atmosphere codes (viz. Martins et al. 2005, Puls et al. 2005).

We are performing a study of Galactic H II regions ionized by only one massive star to check the validity of the ionizing SEDs predicted by the new generation of massive star model atmosphere codes. In forthcoming papers we will present the complete combined study of the various nebulae and their ionizing stars; here we show some preliminary results from our study of M 43.

2. OBSERVATIONAL DATASET

Stellar and nebular observations were carried out with the Wide Field Camera (WFC; narrow-band imaging in H α , H β , [O III] and [S II]) and the Intermediate Resolution Spectrograph (IRS; spectroscopy of the ionizing star in the range 4000-5000 Å + the H α region, and long slit nebular spectroscopy in the range 3600-9700 Å); both instruments attached to the Isaac Newton Telescope (INT). The long slit nebular observations were divided into smaller apertures along the nebular radii.

¹LUTH, Observatoire de Meudon, France.

²Instituto de Astrofísica de Canarias, Spain.

³Instituto de Astronomía, Universidad Nacional Autónoma de México, México.




Fig. 1. Fitting of FASTWIND synthetic profiles (broadened to $v \sin i = 210 \text{ km s}^{-1}$) to the optical H and He lines of HD 37061. Two sets of stellar parameters are shown to illustrate the accuracy of the stellar parameters determination.

3. STELLAR AND NEBULAR ANALYSES

M 43 is an apparently spherical H II region, ionized by HD 37061 (classified as B1 V, though our spectrum clearly shows the He II 4541 line, indicating that the star is rather B0 V - B0.5 V). The stellar parameters of HD 37061 were obtained by visual fitting of FASTWIND (Puls et al. 2005) synthetic profiles to the optical H and He lines (see Figure 1). Once the SED resulting from the FASTWIND model was re-scaled to fit $M_V = 3.5$ (from the stellar photometry and considering a distance of 450 pc), a value of $\log Q(H^0) = 47.2$ could be derived. This later value is in good agreement with the nebular H_α luminosity calculated from the WFC H_α image, indicating that the nebula is ionization bounded. From the nebular [S II] 6731/6716 line ratio we inferred $N_e = 550 \text{ cm}^{-3}$. In a first approach, we have used the nebular abundances derived by Rodríguez (1999)

4. PHOTOIONIZATION MODELS

Figure 2 illustrates three of the diagrams we use to compare the predictions of photoionization models (CLOUDY, Ferland et al. 1998) with the observed nebular constraints. The first one is an indicator of the nebular temperature, while the other two illustrate the nebular ionization structure. These plots compare three constant density spherical CLOUDY models: two of them take the SEDs from FASTWIND models with different stellar parameters, while the third one was calculated using the SED from a TLUSTY (Hubeny & Lanz, 1995) model with the same parameters as the FASTWIND model which best fits




Fig. 2. Spatial variation of various nebular line ratios resulting from three CLOUDY constant density spherical models. Each model uses a different predicted SED. FASTWIND models are the ones used in Figure 1 (same color code); a TLUSTY model with the same stellar parameters as the FASTWIND model which best fits the optical HHe lines (Figure 1) has also been considered for comparison. Nebular observations from long slit spectra are also included.

the optical HHe lines. The differences between the photoionization models are important. Note, however, that geometries other than spherical and non-constant density distributions can affect the result (Morisset et al. 2005). This will be also taken into account when inferring which model better fit the nebular constraints. In forthcoming papers we will present the complete analysis in detail.

REFERENCES

- Ferland, G. J., et al. 1998, *PASP*, 110, 761
- Gabler, R., et al. 1989, *A&A*, 226, 162
- Giveon, U., et al. 2002, *ApJ*, 566, 880
- Hillier, D. J., & Miller, D. L. 1998, *ApJ*, 496, 407
- Hubeny, I., & Lanz, T. 1995, *ApJ*, 439, 875
- Kunze, D., 1994, PhD thesis, Munich, Germany
- Kurucz, R. L. 1992, *IAU Symp.* 149, 225
- Martins, F., Schaerer, D., & Hillier, D. J. 2005, *A&A*, 436, 1049
- Mihalas, D., & Auer, L. H. 1970, *ApJ*, 160, 1161
- Morisset, C., Schaerer, D., Bouret, J.-C., & Martins, F. 2004, *A&A*, 415, 577
- Morisset, C., Stasińska, G., & Peña, M. 2005, *MNRAS*, 360, 499
- Najarro, F., et al. 1996, *A&A*, 306, 892
- Pauldrach, A. W. A., Hoffmann, T. L., & Lennon, M. 2001, *A&A*, 375, 161
- Puls, J., et al. 2005, *A&A*, 435, 669
- Rodríguez, M. 1999, *A&A*, 351, 1075
- Rubin, R. H., Kunze, D., & Yamamoto, T. 1995, *ASP Conf. Ser.* 78: *Astrophysical Applications of Powerful New Databases*, 78, 479
- Sellmaier, F. H., Yamamoto, T., Pauldrach, A. W. A., & Rubin, R. H. 1996, *A&A*, 305, L37
- Stasińska, G., & Schaerer, D. 1997, *A&A*, 322, 615